


23rd WPI IIS Seminar

“Biogenesis and functions of PIWI-interacting RNAs in *Drosophila* germline”

Mobilization of transposable elements (TEs) can lead to natural insertion mutations that generally have negative effects on the host genome. Thus, host species have evolved control mechanisms that restrict TE activity. One such mechanism is RNAi/RNA silencing, where small RNAs of 20–30 nucleotides (nt) in length trigger multiple forms of sequence-specific gene silencing. In the *Drosophila* germline, PIWI-interacting RNAs (piRNAs) are produced from single-stranded long non-coding RNAs that are transcribed from intergenic elements, piRNA clusters, and are loaded onto PIWI proteins to yield piRISCs. However, the underlying mechanisms of how piRNAs are produced and function in TE silencing remain unknown.



Speaker: Dr. Mikiko C. Siomi

*Department of Biophysics and Biochemistry,
Graduate School of Science, the University of Tokyo*

Date: Monday, November 11, 2013

Time: 16:00-17:00

**Venue: Room #402, 4F, Health and Medical Science
Innovation Laboratory, University of Tsukuba**



IIS

INTERNATIONAL INSTITUTE FOR INTEGRATIVE
SLEEP MEDICINE



**Contact: International Institute for Integrative Sleep Medicine
Phone: 029-853-5857 (ext. 5857)**