

Press Release

2022.12.8 | International Institute for Integrative Sleep Medicine (WPI-IIIS)

Unravelling the secrets of a good night's sleep

Researchers from the University of Tsukuba identify a signaling pathway within brain cells that regulates how long and how deeply we sleep

Tsukuba, Japan - A good night's sleep can work wonders for both mind and body. But what is it that determines how much we need to sleep, and what can cause us to sleep more deeply?

In a new study, researchers from the University of Tsukuba have now provided some answers, revealing a signaling pathway within brain cells that regulates the length and depth of sleep.

"We examined genetic mutations in mice and how these affect their patterns of sleep," says senior author of the study, Professor Hiromasa Funato. "We identified a mutation that led to the mice sleeping much longer and more deeply than usual." The researchers found that this was caused by low levels of an enzyme called histone deacetylase 4 (HDAC4), which is known to suppress the expression of target genes.


Previous studies on HDAC4 have shown that it is greatly affected by the attachment of phosphate molecules in a process known as phosphorylation. When this occurs, HDAC4 tends to move away from the cell nucleus, and the suppression of certain proteins is reduced. The researchers were interested in whether this phosphorylation of HDAC4 would affect sleep.

"We focused on a protein called salt-inducible kinase 3, otherwise known as SIK3, which phosphorylates HDAC4," says Professor Funato. "We previously found that this protein has strong effects on sleep." The team found that when there was a lack of SIK3 or when HDAC4 was modified to prevent phosphorylation, the mice slept less. In contrast, when the mice had a more active version of SIK3,

which increased the phosphorylation of HDAC4, they slept a lot more. They also identified a further protein, LKB1, which phosphorylates SIK3, and has similar sleep-suppressing effects when deficient.

“Our findings indicate that there is a signaling pathway within brain cells from LKB1 to SIK3 and then to HDAC4,” says study co-senior author, Professor Masashi Yanagisawa. “This pathway leads to the phosphorylation of HDAC4, which promotes sleep, most probably because it affects the expression of sleep-promoting genes.”

The team carried out further experiments to identify the brain cells in which these pathways regulate sleep. This involved altering the amounts of SIK3 and HDAC4 in different cell types and brain regions. The results indicated that signaling within the cells of the cortex regulates the depth of sleep, while signaling within the hypothalamus regulates the amount of deep sleep. For both brain regions, the excitatory neurons, which can activate other neurons, were identified as playing a key role.



These results provide an important insight into how sleep is regulated, which could potentially lead to a greater understanding of sleep disorders as well as the development of new treatments.

####

The article, “Kinase signalling in excitatory neurons regulates sleep quantity and depth,” was published in *Nature* at DOI: 10.1038/s41586-022-05450-1.

Funding: This work was supported by the World Premier International Research Center Initiative from MEXT, JSPS KAKENHI, JST CREST, AMED, JSPS DC2, University of Tsukuba Basic Research Support Program Type A, Funding Program for World-Leading Innovative R&D on Science and Technology (FIRST Program).

Summary: Researchers from the University of Tsukuba, Japan, have identified a signaling pathway that regulates sleep. By studying genetic mutations in mice, they were able to show that an enzyme called histone deacetylase 4 (HDAC4), which is known to suppress gene expression, plays a key role in regulating the amount and depth of sleep. Further experiments revealed that this regulatory signaling in the excitatory neurons of the cortex and hypothalamus regulates sleep quantity and quality, respectively.

Tweets

Tweet #1: Are you a light sleeper or a heavy sleeper? Signaling pathways within brain cells may help us understand why.

Tweet #2: Signaling pathways within brain cells may help ensure a good night's sleep.

Keywords

Primary Keyword: Life Sciences

Additional Keywords: Sleep, Regulatory mechanisms, Intracellular signaling pathways, Regulation by phosphorylation, Kinase activity, Slow wave sleep, Neurons, Gene expression regulation

Bibliographic information

Staci J. Kim, Noriko Hotta-Hirashima, Fuyuki Asano, Tomohiro Kitazono, Kanako Iwasaki,

Shinya Nakata, Haruna Komiya, Nodoka Asama, Taeko Matsuoka, Tomoyuki Fujiyama, Aya Ikkyu, Miyo Kakizaki, Satomi Kanno, Jinhwan Choi, Deependra Kumar, Takumi Tsukamoto, Asmaa Elhosainy Mohamed Mahmoud, Seiya Mizuno, Shinichi Miyazaki, Yousuke Tsuneoka, Fumihiro Sugiyama, Satoru Takahashi, Yu Hayashi, Masafumi Muratani, Qinghua Liu, Chika Miyoshi, Masashi Yanagisawa, Hiromasa Funato (2022)
“Kinase signalling in excitatory neurons regulates sleep quantity and depth” published in ***Nature*** at DOI: 10.1038/s41586-022-05450-1.

Rui Zhou, Guodong Wang, Qi Li, Fanxi Meng, Can Liu, Rui Gan, Dapeng Ju, Meimei Liao, Junjie Xu, Di Sang, Xue Gao, Shuang Zhou, Kejia Wu, Quanzhi Sun, Ying Guo, Chongyang Wu, Zhiyu Chen, Lin Chen, Bihan Shi, Haiyan Wang, Xia Wang, Huaiye Li, Tao Cai, Bin Li, Fengchao Wang, Hiromasa Funato, Masashi Yanagisawa, Eric Erquan Zhang, Qinghua Liu (2022)
“A signaling pathway for transcriptional regulation of sleep amount in mice” published in ***Nature*** at DOI: 10.1038/s41586-022-05510-6.

Media Contact

Alliance and Communication Unit, International Institute for Integrative Sleep Medicine (WPI-IIIS), University of Tsukuba, Japan
E-mail: wpi-iiis-alliance@ml.cc.tsukuba.ac.jp